



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

AIRS-MLS Upper Tropospheric Water Vapor Comparisons

Eric Fetzer

Jet Propulsion Laboratory, California Institute of Technology

AIRS Science Team Meeting, Greenbelt, MD

27 September 2006



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Executive Summary

AIRS-MLS upper trop. water vapor

- **Excellent agreement ($\sim 30\%$ RMS; $< 5\%$ bias) at 250 hPa for non-polar latitudes.**
- **Poorer agreement at 300, 200 & 150 hPa**
 - *Different sampling distributions.*
 - *MLS $\sim 30\%$ dry at 300 hPa.*
- **Ranked statistics (e. g. medians, percentiles) often agree despite large RMS differences**
 - *Highly non-gaussian data with many outliers*
- **AIRS less sensitive in stratosphere and tropical upper troposphere**
 - *AIRS may have NO skill (except climatology) down to 300 hPa over poles.*
 - *Some sensitivity to 150 hPa in tropics.*
- **MLS appears *more* strongly affected by (ice) clouds than is AIRS.**
 - *Most pronounced in the moist tropics*
 - *Later data versions may fix this.*




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

The Instruments

- **AIRS: Atmospheric Infrared Sounder on Aqua**
 - *Sensitive to ~ 0.1 mm total water (10-20 ppmv in Gettelman et al. 2004, GRL).*
- **MLS: Microwave Limb Sounder on Aura**
 - *Water vapor from 316 hPa upward.*
 - *Sensitive down to very low amounts (a few ppmv).*



The samples are *minutes* apart.



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Some Questions

- **Where do AIRS and MLS have similar water vapor observing characteristics?**
 - *Looked at distributions, ranked statistics, summaries (mean, std. dev.), correlation, linearity.*
- **Where (and why) do they observe differently?**
- **What are the effects of clouds on sampling?**
- **How do these vary between seasons?**
 - *Look at all AIRS-MLS matches for 2005.*



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Use Nearest Neighbor Matching

Why:

- The goal of this work is reconciling the two data sets
 - *~25% RMS & ~5% bias are 'close enough'*
- *Sampling effects of clouds are critical to understanding climatologies*
 - *Both instrument flag 'undesirable' scenes. Keep track of these...*



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

More Matching Issues...

- **Count only AIRS-MLS match-ups.**
 - ***This under-represents AIRS sampling by a factor of ~100.***
- **Place both data sets on the AIRS standard levels of 300, 250, 200 and 150 hPa.**
 - ***AIRS: geometric mean of layers.***
 - ***MLS: Log(mixing ratio) linear in Log(p).***



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Quality Flagging

- **Both instruments use quality flags**
 - **AIRS**
 - Scattering of microwaves by precipitation, or cloud cover greater than 50-70%.
 - *Use Qual_Temp_Profile_Mid = 0.*
 - **MLS**
 - Microwave scatter from ice particles larger than ~10 microns.
 - *Use Quality >5.0 at 316 hPa, >0.3 above*



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

First Lesson: Important to exclude AIRS water vapor above tropopause *AIRS versus MLS at 200 hPa*

Exclude these:

South Polar,
above AIRS
tropopause

North Polar,
above AIRS
tropopause
(agree in the
mean).



Keep these

Non-polar,
beneath
AIRS
tropopause




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Sampling by month and latitude

- **Examine:**
 - *Twelve months in 2005.*
 - *Twelve 15-degree latitude bands.*
 - *Four pressures: 300, 250, 200 & 150 hPa.*
- **We see:**
 - *AIRS is often 'stratospheric' down to 300 hPa over poles.*
 - *Many familiar regions of poorer AIRS yields:*
 - Subtropical stratus.
 - Midlatitude storms.
 - Polar regions in summer.
 - *Both AIRS and MLS have low yields in regions of deep convection*
 - Very important for MLS ice-water vapor climatologies.




National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Tropics, 250 hPa

Small biases, RMS agreement to ~30% for all months

Relative differences roughly constant with amount





National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Tropics, 300 hPa

Agreement *poorer* than at 250 hPa

MLS ~30% drier

Relative differences vary with amount






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Tropics, 200 & 150 hPa Differences vary with amount

**NOTE: Sensitivity
threshold varies
with height!**

**Gettelman et al., 2004,
GRL say it's constant
at 10-20 ppmv.**





National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Conclusions and Future Work

- **Agreement to a few percent in mean, 25% RMS at 250 hPa**
 - *consistent with MLS-CFH sonde results by Holger Vömel, Costa Rica.*
- **MLS dry bias of ~30% at 300 hPa noted by Vömel, others**
 - *Tobin shows AIRS dry bias of 10%.*
- **Mixed results at 200 & 150 hPa**
 - *Low-end insensitivity by AIRS could explain this.*
- **Examining effects of cloud sampling**
 - *Complementary data sets in tropics*
 - MLS misses much water vapor -- but samples ice!
 - AIRS nicely samples water vapor.
- **Manuscript(s) currently in preparation.**